

DTIC FILE COPY

2

AD-A228 854

LINEAR-TIME SEPARATION ALGORITHMS  
FOR THE THREE-INDEX  
ASSIGNMENT POLYTOPE

Egon Balas  
Carnegie Mellon University

and

Liqun Qi  
University of New South Wales


Carnegie Mellon University

PITTSBURGH, PENNSYLVANIA 15213

DTIC  
SELECTED  
NOV 08 1990  
S E

GRADUATE SCHOOL OF INDUSTRIAL ADMINISTRATION

WILLIAM LARIMER MELLON, FOUNDER



DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

90-11-7-005

**LINEAR-TIME SEPARATION ALGORITHMS  
FOR THE THREE-INDEX  
ASSIGNMENT POLYTOPE**

Egon Balas  
Carnegie Mellon University

*and*

Liqun Qi  
University of New South Wales

September 1990

**SDTIC  
ELECTE  
NOV 08 1990  
E D**

The research underlying this report was supported by the National Science Foundation through Grant No. DDM-8901495 and the Office of Naval Research through Contract N00014-85-K-0198. Also partially supported by the Australian Research Council.

Management Science Research Group  
Graduate School of Industrial Administration  
Carnegie Mellon University  
Pittsburgh, PA 15213

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

# Abstract

Balas and Saltzman [1] identified several classes of facet inducing inequalities for the three-index assignment polytope, and gave  $O(n^4)$  separation algorithms for two of them. We give  $O(n^3)$  separation algorithms for these two classes of facets, and also for a third class. Since the three-index assignment problem has  $n^3$  variables, these algorithms are linear-time and their complexity is best possible.

## Keywords:

Three-index assignment  
Facets  
Cliques  
Odd holes



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Dist. "A" per telecon Dr. Donald Wagner.  
Office of Naval Research/code 1111MA.

VHC

11/08/90

## 1. Introduction

Consider three disjoint  $n$ -sets and the collection of all weighted triplets with one element in each  $n$ -set. The three-index assignment (or three-dimensional matching) problem asks for a minimum-weight set of triplets that partitions the union of the three  $n$ -sets. It can be stated as the following 0-1 programming problem:

$$\begin{aligned}
 \min \quad & \sum \{c_{ijk}x_{ijk} : i \in I, j \in J, k \in K\}, \\
 \text{s.t.} \quad & \sum \{x_{ijk} : j \in J, k \in K\} = 1, \quad \forall i \in I, \\
 & \sum \{x_{ijk} : i \in I, k \in K\} = 1, \quad \forall j \in J, \\
 & \sum \{x_{ijk} : i \in I, j \in J\} = 1, \quad \forall k \in K, \\
 & x_{ijk} \in \{0, 1\}, \quad \forall i, j, k,
 \end{aligned} \tag{1.1}$$

where  $I, J$  and  $K$  are disjoint sets with  $|I| = |J| = |K| = n$ . Let  $A$  be the coefficient matrix of the constraint set of (1.1). Then  $R = I \cup J \cup K$  is the row index set of  $A$ . Let  $S$  be the column index set of  $A$ . Let  $G_A$  be the intersection graph of  $A$ . Then  $S$  is the node set of  $G_A$ . Let

$$P = \{x \in \mathbb{R}^{n^3} : Ax = e, x \geq 0\},$$

where  $e = (1, \dots, 1)^T \in \mathbb{R}^{3n}$ . Then

$$P_I = \text{conv}\{x \in \{0, 1\}^{n^3} : x \in P\}$$

is the three-index assignment polytope of order  $n$ .

Balas and Saltzman [1] described the cliques of  $G_A$  as belonging to three distinct classes and showed that cliques in two of the three classes induce inequalities that define facets of  $P_I$ . Furthermore, they gave an  $O(n^4)$  procedure for finding a facet-defining clique-inequality violated by a given noninteger  $x \in P$ , or showing that no such inequality exists. They also described in [1] the odd holes of  $G_A$  and identified two classes of facets associated with odd holes that are easy to generate. One class has coefficients of 0 and 1, the other class coefficients of 0, 1 and 2.

In Section 2 of this paper, we give two  $O(n^3)$  separation algorithms for the above mentioned two classes of facet defining clique-inequalities. In Section 3, we give an  $O(n^3)$  separation algorithm for facet inducing inequalities attributed with lifted odd holes of length 5, having coefficients equal to 0 or 1. Since the number of variables of the three-index assignment problem is  $n^3$ , the complexity of these procedures is of the lowest possible order. Our procedures can be used to speed up the performance of the branch and bound algorithm

for the three-index assignment problem developed in [2], which uses facet inducing inequalities as ingredients of a Lagrangian relaxation to obtain strong lower bounds. Our procedures can be extended to more general cases. In Section 4, we present an  $O(n^{h+1})$  procedure for detecting a violated facet-defining inequality of  $P_I$ , which is associated with a  $2h + 1$ -hole of  $G_A$  and belongs to a subclass of the first odd-hole-associated class of facets of [1].

We use the same notation as [1]. Especially,  $a^s$  means a column of  $A$  associated with  $s \in S$ . We may specify  $s = \{i, j, k\}$  for  $i \in I, j \in J, k \in K$ . For a set  $Q \subseteq R$ , we use  $Q_I, Q_J$  and  $Q_K$  to denote the parts of  $Q$  in  $I, J$  and  $K$  respectively.

For literature on the three-index assignment problem, also see [3], ..., [8].

## 2. Facets Induced by Cliques

For every  $s \in S$ , let

$$C(s) = \{t \in S : a^s \cdot a^t \geq 2\}.$$

Then the node set  $C(s)$  induces a clique of size  $3n - 2$  in  $G_A$  [1, Proposition 2.2]. Such a clique is called a clique of class 2. Furthermore, for  $n \geq 3$ , the inequality

$$x(C(s)) \equiv \sum \{x_t : t \in C(s)\} \leq 1 \quad (2.1)$$

defines a facet of  $P_I$  for every  $s \in S$  [1, Theorem 3.3]. Given a noninteger  $x \in P$ , an  $O(n^4)$  procedure was presented in [1, Section 4] to detect whether any inequality induced by a clique of class 2 is violated. We now present an  $O(n^3)$  procedure to do this.

**Algorithm 2.1.** Suppose that  $x$  is a noninteger point in  $P$ . Let  $\nu$  be an integer greater than or equal to 4.

*Step 1.* Let  $d_s = 0$  for all  $s \in S$ .

*Step 2.* Check  $x_t$  for all  $t \in S$ . If

$$x_t \geq \frac{1}{\nu n}, \quad (2.2)$$

then set  $d_s := d_s + x_t$  for all  $s \in C(t)$ . If  $d_s > 1$ , then stop: (2.1) is violated by  $x$  with this  $s$ . Otherwise, continue.

*Step 3.* For  $s \in S$ , if

$$d_s > \frac{\nu - 3}{\nu}, \quad (2.3)$$

then check whether the inequality (2.1) associated with  $s$  is violated and if so, then stop. Else continue. ■

**Theorem 2.2.** *Algorithm 2.1 determines in  $O(n^3)$  steps whether a given  $x \in P$  violates a facet defining inequality induced by a clique of class 2. The value of  $\nu$  which minimizes the complexity of the algorithm is 6.*

To prove this theorem we need two lemmas. The first lemma is essentially Lemma 4.1 of [1]. We repeat it here for convenience:

**Lemma 2.3.** *For any  $x \in P$  and any positive number  $\lambda$ , the number of components with value  $\geq \lambda$  is  $\leq n/\lambda$ . ■*

Lemma 4.1 of [1] states this for  $1/\lambda$  integer, but the result holds for reals as well. The proof in [1] is somewhat indirect. In fact, the sum of components of  $x \in P$  is always  $n$ , and all the components of  $x$  are nonnegative. The conclusion thus follows.

**Lemma 2.4.** *For any  $x \in P$  and any positive number  $\lambda$ , the number of  $s \in S$  such that  $x(C(s)) \geq \lambda$  is  $\leq n(3n-2)/\lambda$ .*

*Proof:* Consider

$$\sum \{x(C(s)) : s \in S\}.$$

For each  $t \in S$ , the number of distinct  $s$  such that  $t \in C(s)$  is  $3n-2$ . Thus,

$$\sum \{x(C(s)) : s \in S\} = (3n-2) \sum \{x_t : t \in S\} = (3n-2)n.$$

Since  $x(C(s)) \geq 0$  for each  $s$ , the conclusion follows. ■

*Proof of Theorem 2.2:* We first show that Algorithm 2.1 works. By Proposition 2.2 of [1],  $|C(s)| = 3n-2$  for every  $s \in S$ . Suppose that (2.1) is violated for an  $s \in S$ . Then

$$d_s := \sum \{x_t : t \in C(s), x_t \geq \frac{1}{\nu n}\}$$

$$> 1 - \sum \{x_t : t \in C(s), x_t < \frac{1}{\nu n}\}$$

$$\geq 1 - \frac{3n-2}{\nu n} \geq \frac{\nu-3}{\nu},$$

i.e., (2.3) holds and the violation is discovered in Step 3. Therefore, Algorithm 2.1 determines whether a given  $x \in P$  violates a facet defining inequality induced by a clique of class 2.

We now consider the complexity of the algorithm. The complexity of Step 1 is  $n^3$ . According to Lemma 2.3, there are at most  $\nu n^2$  components of  $x$  satisfying (2.2). For each  $t$ , the number of  $s$  in  $C(t)$  is  $3n - 2$ . Hence, the complexity of Step 2 is  $\nu n^2(3n - 2)$ . Finally, if (2.3) holds, then

$$x(C(s)) \geq d_s \equiv \sum \{x_t : t \in C(s), x_t \geq \frac{1}{\nu n}\} > \frac{\nu - 3}{\nu}.$$

By Lemma 2.4, the number of  $s$  such that (2.3) holds is not greater than  $\frac{\nu n(3n-2)}{\nu-3}$ . The number of basic operations to check (2.1) is  $3n - 2$ . We also need to use  $n^3$  basic steps to check (2.3). Therefore, the complexity of Step 3 is  $n^3 + \frac{\nu n(3n-2)^2}{\nu-3}$ . Hence, the overall complexity of Algorithm 2.1 is  $O(n^3)$ . The value of  $\nu$  that minimizes the maximum of the two polynomials expressing the complexity of Steps 2 and 3 is easily seen to be 6. ■

We now consider facets induced by cliques of class 3 [1]. Suppose that  $s, t \in S$  such that

$$a^s \cdot a^t = 0. \quad (2.4)$$

Let  $s = (i_s, j_s, k_s)$ ,  $t = (i_t, j_t, k_t)$ ,  $t_1 = (i_s, j_t, k_t)$ ,  $t_2 = (i_t, j_s, k_t)$ ,  $t_3 = (i_t, j_t, k_s)$ , and

$$C(s, t) = \{s, t_1, t_2, t_3\}.$$

Then the node set  $C(s, t)$  induces a (4-)clique in  $G_A$  [1, Proposition 2.3]. Such a clique is called a clique of class 3. The number of cliques of class 3 is  $\frac{1}{4}n^3(n - 1)^3$  [1, Corollary 2.5]. Furthermore, for  $n \geq 4$ , the inequality

$$x(C(s, t)) \equiv \sum \{x_t : t \in C(s, t)\} \leq 1 \quad (2.5)$$

defines a facet of  $P_I$  for every  $s, t \in S$  satisfying (2.4) [1, Theorem 3.7]. Given a noninteger  $x \in P$ , an  $O(n^4)$  procedure was also presented in [1, Section 4] to detect whether any inequality induced by a clique of class 3 is violated. We now present an  $O(n^3)$  procedure to do this.

**Algorithm 2.5.** Suppose that  $x$  is a noninteger point in  $P$ . Check  $x_s$  for  $s \in S$ . If

$$\frac{1}{4} < x_s < 1, \quad (2.6)$$

then check  $x_p$  for  $p \in S$  satisfying

$$a^s \cdot a^p = 1. \quad (2.7)$$

If

$$x_p > \frac{1 - x_s}{3}, \quad (2.8)$$

then for  $t \in S$  such that  $a^t \cdot a^s = 0$  and

$$a^t \cdot a^p = 2, \quad (2.9)$$

check whether (2.5) is violated or not. ■

**Theorem 2.6.** *Algorithm 2.5 determines in  $O(n^3)$  steps whether a given  $x \in P$  violates a facet defining inequality induced by a clique of class 3.*

*Proof:* We first show that Algorithm 2.5 works. Let  $C(s, t)$  be the node set of a clique of class 3. Since  $|C(s, t)| = 4$ , if  $x \in P$  violates (2.5), then  $x$  has at least one component  $> \frac{1}{4}$ . By Proposition 2.4 of [1], there is no loss of generality in assuming that this happens for the components indexed by  $s$ , i.e., that  $x_s > \frac{1}{4}$ . Since  $x \in P$ , we have

$$\sum \{x_{i,j,k} : j \in J, k \in K\} = 1, \quad (2.10)$$

$$\sum \{x_{i,j,k} : i \in I, k \in K\} = 1, \quad (2.11)$$

$$\sum \{x_{i,j,k} : i \in I, j \in J\} = 1. \quad (2.12)$$

If  $x_s = 1$ , then  $x_p = 0$  for any  $p$  such that  $a^p \cdot a^s = 1$ , by (2.10), (2.11) and (2.12). In this case, for any  $t$  such that  $a^t \cdot a^s = 0$ , the left hand side of (2.5) is 1. Thus, if  $x \in P$  violates (2.5),  $x_s < 1$ . This justifies (2.6). Since  $|C(s, t)| = 4$ , if (2.5) is violated, then (2.8) must hold for at least one  $p \in C(s, t)$ ,  $p \neq s$ . Therefore, this algorithm determines whether a given  $x \in P$  violates a facet defining inequality induced by a clique of class 3.

We now consider the complexity issue. The complexity to check all  $s$  is  $O(n^3)$ . According to Lemma 2.3, there are at most  $4n$  components of  $x$  satisfying (2.6). For each such component  $x_s$ , there are  $(n-1)^2$  components  $x_p$  satisfying (2.7). So the complexity to find all pairs  $s, p \in S$  satisfying (2.6) and (2.7) is  $O(n^3)$ . By (2.10), (2.11) and (2.12), for given  $x_s$  satisfying (2.6), there are at most 6  $p$  such that (2.7) and (2.8) hold. When  $s$  and  $p$  are fixed and (2.7) is satisfied, there are  $n-1$   $t$  satisfying (2.4) and (2.9). One needs 3 additions and one comparison to check (2.5). Therefore, the complexity of the remaining part of the algorithm is  $O(n^2)$ . Hence, the overall complexity of Algorithm 2.5 is  $O(n^3)$ . ■

In the proof, we see that the main work is to check all  $s \in S$ . This work cannot be reduced. So this order cannot be further improved. Similarly, the order in Theorem 2.2 is also the minimum.

By [1], only these two classes of cliques induce facets of  $P_I$ . Combining Theorems 2.2 and 2.6, we have

**Theorem 2.7.** *One can determine in  $O(n^3)$  steps whether a given  $x \in P$  violates a facet defining inequality induced by a clique of  $G_A$ . ■*



### 3. Facets Associated with Five-Holes

Assume that  $n \geq 3$ . Let  $Q \subseteq R$ ,  $|Q| = 2h + 1$  for some positive integer  $h$  with  $h \leq n - 1$ ,  $1 \leq |Q_L| \leq h$ ,  $L = I, J, K$ , and let

$$S(Q) := \{s \in S : \sum \{a'_q : q \in Q\} \geq 2\}. \quad (3.1)$$

Then the inequality

$$\sum \{x_s : s \in S(Q)\} \leq h \quad (3.2)$$

defines a facet of  $P_I$  [1, Theorem 6.1]. These facets can be regarded as lifted from odd hole inequalities. The number of distinct inequalities (3.2) is  $O(2^{3n})$  [1, Proposition 6.5]. In particular, facet defining inequalities (2.1) are special cases of (3.2) with  $h = 1$ .

Denote the set of  $Q$  satisfying the above requirements with a fixed  $h$  by  $S^h$ . Let  $\Phi_h = \{S(Q) : Q \in S^h\}$ , where  $S(Q)$  is defined by (3.1). Thus, Algorithm 2.1 is an  $O(n^3)$  algorithm to detect violated facets induced by sets in  $\Phi_1$ . In this section, we present an  $O(n^3)$  procedure to detect whether there exists a facet induced by a set in  $\Phi_2$ , which is violated by a given noninteger  $x \in P$ . Notice that the number of facets defined by (3.2) with  $h = 2$  is  $O(n^5)$ . Without loss of generality, we may assume that such an  $x$  does not violate any inequality (2.1), i.e., assume that Algorithm 2.1 was applied to  $x$  first.

Suppose that  $x \in P$ . For  $i \in I, j \in J, k \in K$ , we will denote

$$x(i, j, K) := \sum \{x_{ijk'} : k' \in K\}, \quad (3.3)$$

$$x(i, J, k) := \sum \{x_{ij'k} : j' \in J\}, \quad (3.4)$$

$$x(I, j, k) := \sum \{x_{i'jk} : i' \in I\} \quad (3.5)$$

In the following, we establish some formulas to calculate  $x(S(Q))$  for  $Q \in S^2$ . We consider two cases.

First, suppose that  $s, t \in S$  satisfy

$$a^s \cdot a^t = 1.$$

Then

$$Q \equiv s \cup t \in S^2.$$

Assume that  $s = (i_s, j_s, k_s), t = (i_t, j_t, k_t)$ . It is not difficult to see that if  $k_s = k_t$ , then

$$x(S(Q)) = x(C(s)) + x(C(t)) + x(i_s, j_t, K) + x(i_t, j_s, K) - x_{i_s, j_t, k_s} - x_{i_t, j_s, k_s}; \quad (3.6)$$

if  $j_s = j_t$ , then

$$x(S(Q)) = x(C(s)) + x(C(t)) + x(i_s, J, k_t) + x(i_t, J, k_s) - x_{i_s, j_t, k_t} - x_{i_t, j_s, k_s}; \quad (3.7)$$

and if  $i_s = i_t$ , then

$$x(S(Q)) = x(C(s)) + x(C(t)) + x(I, j_s, k_t) + x(I, j_t, k_s) - x_{i_s, j_s, k_t} - x_{i_t, j_t, k_s}. \quad (3.8)$$

Next, suppose that  $s, t \in S$  satisfy

$$a^s \cdot a^t = 2. \quad (3.9)$$

Again assume that  $s = (i_s, j_s, k_s), t = (i_t, j_t, k_t)$ . Then define for the ordered pair  $(s, t)$ ,

$$x(s, t) := x(i_t, J, k_t) + x(i_t, j_t, K) - 2x_{i_t, j_t, k_t}, \quad (3.10)$$

if  $i_s \neq i_t$ ;

$$x(s, t) := x(i_t, j_t, K) + x(I, j_t, k_t) - 2x_{i_t, j_t, k_t}, \quad (3.11)$$

if  $j_s \neq j_t$ ;

$$x(s, t) := x(I, j_t, k_t) + x(i_t, J, k_t) - 2x_{i_t, j_t, k_t}, \quad (3.12)$$

if  $k_s \neq k_t$ . Notice that in (3.10)  $j_t = j_s$  and  $k_t = k_s$ , in (3.11)  $i_t = i_s$  and  $k_t = k_s$ , and in (3.12)  $i_t = i_s$  and  $j_t = j_s$ . Thus, (3.10) can be written with  $j_s$  and  $k_s$  substituted for  $j_t$  and  $k_t$ , and there are similar expressions for (3.11) and (3.12).

In the case that (3.9) holds, we may expand  $s \cup t$  to a set  $Q \in S^2$  by adding an adequate element. For example, if  $i_s \neq i_t$ , we may add  $j \neq j_s$  and let

$$Q_I = \{i_s, i_t\}, Q_J = \{j_s, j\}, Q_K = \{k_s\}, \quad (3.13)$$

or we may add  $k \neq k_s$  and let

$$Q_I = \{i_s, i_t\}, Q_J = \{j_s\}, Q_K = \{k_s, k\}. \quad (3.14)$$

Notice that there are  $2(n-1)$  ways to expand a given pair of  $s$  and  $t$  satisfying (3.9) to a set  $Q \in S^2$ . We may also calculate  $x(S(Q))$  by expressions similar to (3.6)-(3.8). For example, in case (3.13), one can check that

$$x(S(Q)) = x(C(s)) + x(s, t) + x(i_s, j, K) + x(i_t, j, K) + x(I, j, k_s) - 2x_{i_s, j, k_s} - 2x_{i_t, j, k_s}, \quad (3.15)$$

where  $x(s, t)$  is given by (3.10). Note that if  $i_s \neq i_t$ , there are two possibilities, namely, (3.13) and (3.14). Therefore, there are six cases totally, i.e., two cases for  $i_s \neq i_t$ , two cases for  $j_s \neq j_t$ , and two cases for  $k_s \neq k_t$ . In each of these six cases, there is an expression of type (3.15) to calculate  $x(S(Q))$ . Later, we will refer to these as "expressions of type (3.15)".

We are now ready to describe our algorithm:

**Algorithm 3.1.** Suppose that  $x$  is a noninteger point in  $P$ . Suppose (2.1) holds for all  $s \in S$ .

*Step 1.* Let  $d_s = 0$  for all  $s \in S$ .

*Step 2.* Check  $x_t$  for all  $t \in S$ . If

$$x_t \geq \frac{1}{12n}, \quad (3.16)$$

then let  $d_s := d_s + x_t$  for all  $s \in S$  satisfying

$$a_s \cdot a_t \geq 2.$$

*Step 3.* For all  $s \in S$ , if

$$d_s \geq \frac{1}{12}, \quad (3.17)$$

then set  $d_s := x(C(s))$ .

*Step 4.* For all  $i \in I, j \in J, k \in K$ , calculate  $x(i, j, K)$ ,  $x(i, J, k)$  and  $x(I, j, k)$  by (3.3), (3.4) and (3.5).

*Step 5.* For all  $i \in I, j \in J, k \in K$ , form the sets

$$L(i, J, K) := \{s \in S : d_s > \frac{1}{3}, i_s = i\},$$

$$L(I, j, K) := \{s \in S : d_s > \frac{1}{3}, j_s = j\}$$

and

$$L(I, J, k) := \{s \in S : d_s > \frac{1}{3}, k_s = k\}$$

Store them.

*Step 6.* For each  $s = (i_s, j_s, k_s) \in S$  such that  $d_s > \frac{1}{2}$ , check  $d_t$  for each  $t \in L(s)$ , where

$$L(s) := (L(i_s, J, K) \cup L(I, j_s, K) \cup L(I, J, k_s)) \setminus \{s\}. \quad (3.18)$$

If  $a^t \cdot a^s = 1$  and  $d_t > (2 - d_s)/3$ , calculate  $x(S(Q))$  by the appropriate expression (3.6), (3.7) or (3.8), with  $d_p$  substituted for  $x(C(p))$ ,  $p = s, t$ .

If  $a^t \cdot a^s = 2$  and  $x(s, t) > (2 - d_s)/3$ , calculate  $x(S(Q))$  by the appropriate expression of type (3.15), with  $d_p$  substituted for  $x(C(p))$ ,  $p = s, t$ , for every  $Q \in S^2$  such that  $s \cup t \subseteq Q$ .

If  $x(S(Q)) > 2$  for some  $Q$ , stop: the corresponding inequality (3.2) is violated by  $x$ . Otherwise, continue. ■

**Theorem 3.2.** *Algorithm 3.1 determines in  $O(n^3)$  steps whether a given  $x \in P$  violates a facet defining inequality (3.2) with  $h = 2$ .*

We prove several auxiliary results.

**Lemma 3.3.** *After Step 3 of Algorithm 3.1, for all  $s \in S$ ,*

$$d_s \leq x(C(s)). \quad (3.19)$$

*If  $x(C(s)) > \frac{1}{3}$ , then  $d_s = x(C(s))$ .*

*Proof:* It is obvious that (3.19) holds. Suppose that  $x(C(s)) > \frac{1}{3}$ . Then

$$\begin{aligned} d_s &= \sum \{x_t : t \in C(s), x_t \geq \frac{1}{12n}\} > \frac{1}{3} - \sum \{x_t : t \in C(s), x_t < \frac{1}{12n}\} \\ &\geq \frac{1}{3} - \frac{3n-2}{12n} \geq \frac{1}{12}, \end{aligned}$$

i.e.,  $d_s \geq \frac{1}{12}$  holds and in this case  $d_s$  is set equal to  $x(C(t))$  in Step 3. ■

**Lemma 3.4.** *At Step 6,*

$$|L(s)| \leq 27n, \quad (3.20)$$

$$|\{t \in L(s) : a^s \cdot a^t = 2, x(s, t) > \frac{1}{3}\}| \leq 18, \quad (3.21)$$

where  $L(s)$  is defined by (3.18) and  $x(s, t)$  is defined by (3.10), (3.11) and (3.12).

*Proof:* Suppose that  $s = (i_s, j_s, k_s)$ .

On the one hand, by Lemma 3.3 and the definition of  $L(i_s, J, K)$  at Step 5,

$$\sum \{x(C(t)) : t \in L(i_s, J, K)\} \geq \frac{1}{3} |L(i_s, J, K)|.$$

On the other hand,

$$\begin{aligned} &\sum \{x(C(t)) : t \in L(i_s, J, K)\} \\ &= \sum_{j_t, k_t} \{x(C(t)) : t = (i_s, j_t, k_t)\} \\ &\leq \sum_{j_t, k_t} [\sum_i x_{ij_t k_t} + \sum_j x_{i_s j k_t} + \sum_k x_{i_s j_t k}] \\ &= \sum_i \sum_{j_t, k_t} x_{ij_t k_t} + \sum_{j_t} \sum_{j, k_t} x_{i_s j k_t} + \sum_{k_t} \sum_{j_t, k} x_{i_s j_t k} \end{aligned}$$

$$= \sum_i 1 + \sum_{j_i} 1 + \sum_{k_i} 1 = 3n.$$

Thus,  $|L(i_s, J, K)| \leq 9n$ . Similarly, we can show that  $|L(I, j_s, K)| \leq 9n$  and  $|L(I, j_s, K)| \leq 9n$ . From the definition of  $L(s)$ , we then have (3.20).

It is also seen that

$$\begin{aligned} & \sum \{x(s, t) : t \in S, a^s \cdot a^t = 2\} \\ & \leq 2 \sum_{j, k} x_{i, j, k} + 2 \sum_{i, k} x_{i, j, k} + 2 \sum_{i, j} x_{i, j, k} = 6. \end{aligned}$$

Hence, (3.21) holds. ■

**Lemma 3.5.** Suppose that  $Q \in S^2$ . Let

$$H(Q) := \{s \in S : \sum \{a_q^s : q \in Q\} = 3\}.$$

If the inequality (3.2) with  $h = 2$  is violated by  $x$ , then there are  $s, t \in H(Q)$ ,  $s \neq t$  such that

- (i).  $x(C(s)) > \frac{1}{2}$ ;
- (ii). either  $a^s \cdot a^t = 1$  and  $x(C(t)) > (2 - x(C(s)))/3$ , or  $a^s \cdot a^t = 2$  and  $x(s, t) > (2 - x(C(s)))/3$ .

*Proof:* Without loss of generality, assume that  $Q_I = \{i_1, i_2\}$ ,  $Q_J = \{j_1, j_2\}$ ,  $Q_K = \{k\}$ . Then  $H(Q) = \{(i_1, j_1, k), (i_1, j_2, k), (i_2, j_1, k), (i_2, j_2, k)\}$ , and

$$2 < x(S(Q)) \leq x(C((i_1, j_1, k))) + x(C((i_1, j_2, k))) + x(C((i_2, j_1, k))) + x(C((i_2, j_2, k))).$$

Then at least one term on the right hand side of the above inequality is greater than  $\frac{1}{2}$ . Without loss of generality, assume it is the first term. Let  $s = (i_1, j_1, k)$ . Then (i) holds.

Denote  $t_1 = (i_2, j_2, k)$ ,  $t_2 = (i_1, j_2, k)$ ,  $t_3 = (i_2, j_1, k)$ . Then

$$\begin{aligned} & 2 < x(S(Q)) \\ & \leq x(C(s)) + x(C(t_1)) + x(i_1, j_2, K) + x(i_2, j_1, K) - x_{i_1 j_2 k} - x_{i_2 j_1 k}, \quad (\text{by (3.6)}) \\ & \leq x(C(s)) + x(C(t_1)) + x(s, t_2) + x(s, t_3). \quad (\text{by (3.11) and (3.10)}) \end{aligned}$$

Notice that  $a^s \cdot a^{t_1}$  and  $a^s \cdot a^{t_3} = a^s \cdot a^{t_3} = 2$ . Thus,

$$2 - x(C(s)) < x(C(t_1)) + x(s, t_2) + x(s, t_3).$$

At least one term at the right hand side of above inequality is greater than one third of the left hand side of this inequality. Hence, (ii) holds. ■

*Proof of Theorem 3.2:* By Lemmas 3.5 and 3.3, if the inequality (3.2) with  $h = 2$  is violated by  $x$  for some  $Q$ , there are  $s, t \in S$  such that  $d_s > \frac{1}{2}$ , and either  $a^s \cdot a^t = 1$  and  $d_t > (2 - d_s)/3$ , or  $a^s \cdot a^t = 2$  and  $x(s, t) > (2 - d_s)/3$ .

Thus if  $t \in L(s)$ , i.e., if  $d_t > \frac{1}{3}$ , then Step 6 will detect the violation. To see that this is the case, notice that if  $a^s \cdot a^t = 1$ , then

$$d_t > (2 - d_s)/3 = (2 - x(C(s)))/3 \geq \frac{1}{3},$$

(since  $x(C(s)) \leq 1$  for all  $s \in S$ ), and if  $a^s \cdot a^t = 2$ , then

$$\begin{aligned} d_t = x(C(t)) &\geq x(s, t) \quad (\text{from the definitions}) \\ &> (2 - d_s)/3 = (2 - x(C(s)))/3 \geq \frac{1}{3}, \end{aligned}$$

Thus in both cases  $t \in L(s)$ .

Consider now the complexity of Algorithm 3.1. Steps 1 and 2 are obviously  $O(n^3)$ . In Step 3, the number of  $s \in S$  such that  $d_s \geq \frac{1}{12}$  is  $\leq 12n(3n - 2)$  (From Lemma 2.4 and the fact that  $d_s \leq x(C(s))$ ; and for each number  $s$ , calculating  $x(C(s))$  requires  $3n - 3$  additions; thus the complexity of Step 3 is also  $O(n^3)$ . Further, each of Steps 4 and 5 again requires  $O(n^3)$  operations.

Finally, consider the complexity of Step 6. By Lemmas 3.3 and 2.4, the number of  $s$  such that  $d_s > \frac{1}{2}$  is  $O(n^2)$ . From Lemma 3.4, the cardinality of  $L(s)$  is  $O(n)$ . Forming  $L(s)$  itself is  $O(1)$ , since its constituent sets have been calculated and stored in Step 5. Thus, checking  $d_t$  for each  $t \in L(s)$  for all  $s$  such that  $d_s > \frac{1}{2}$  is  $O(n^3)$ . If  $a^s \cdot a^t = 1$  and  $d_t > (2 - d_s)/3$ , calculating  $x(S(Q))$  is  $O(1)$ , since the expressions  $x(C(s))$ ,  $x(C(t))$  and  $x(i, j, K)$ ,  $x(i, j, k)$ ,  $x(I, j, k)$  have been calculated in Step 3 and Step 4 respectively. On the other hand, from Lemma 3.4 the number of those  $t \in L(s)$  (for given  $s$ ) such that  $a^s \cdot a^t = 2$  and  $x(s, t) > \frac{1}{3}$  is  $O(1)$ . For each such  $t$ , there are  $2(n - 1)$  ways to expand the pair  $(s, t)$  to a set  $Q \in S^2$ , and for each such set  $Q$ , calculating  $x(S(Q))$  is again  $O(1)$ , since the constituent terms of the expressions of type (3.15) needed for this have already been calculated. Thus, the complexity of dealing with all pairs  $(s, t)$  satisfying  $a^s \cdot a^t = 2$  is again  $O(n^3)$ , which leaves the total complexity of Step 6 at  $O(n^3)$ . ■

#### 4. Facets Associated with Other Odd Holes

It is somewhat more complicated to detect violated facets defined by (3.2) with  $h > 2$ . We first discuss some subclasses of  $S^h$  and  $\Phi_h$ . Assume that  $n \geq 3$ . Let  $\tilde{S}^h$  be the subcollection of  $S^h$  such that for any

$Q \in \bar{S}^h$  there exists an  $L = I, J$ , or  $K$  such that  $|Q_L| = 1$ . Note that in this case  $|Q_{L'}| = h$  for  $L' \in \{I, J, K\}$ ,  $L' \neq L$ . Let  $\bar{\Phi}_h = \{S(Q) : Q \in \bar{S}^h\}$ . Clearly,  $\bar{S}^1 = S^1$ ,  $\bar{S}^2 = S^2$ , but  $\bar{S}^h$  is a proper subset of  $S^h$  for  $h > 2$ . Assume that  $h \geq 3$ . We will show that

**Theorem 4.1.** *For a given  $x \in P$ , it is possible in  $O(n^{h+1})$  steps to detect whether there exists a facet of  $P_I$ , which is defined by (3.2) with  $Q \in \bar{S}^h$  and violated by  $x$ .*

Again, we prove several auxiliary results.

Suppose that  $x \in P$  does not violate any inequality

$$x(S(Q)) \leq r. \quad (4.1)$$

with  $Q \in \bar{S}^r$ ,  $r = 1, 2, \dots, h-1$ , but violates (4.1) with  $h = r$  and a set  $Q \in \bar{S}^h$ . Suppose that  $h$  is fixed, not dependent on  $n$ .

**Lemma 4.2.** *Let  $H(Q)$  be defined as in Lemma 3.5. Then there exists an  $s \in H(Q)$  such that*

$$x(C(s)) > \frac{1}{h}. \quad (4.2)$$

*Proof:* Clearly,

$$h < x(S(Q)) \leq \sum \{x(C(s)) : s \in H(Q)\}.$$

However, for  $Q \in \bar{S}^h$ ,  $|H(Q)| \leq h^2$ . The conclusion thus follows. ■

Since  $Q \in \bar{S}^h$ , without loss of generality, assume that  $Q_K = \{k\}$ . Then  $|Q_I| = |Q_J| = h$ . Suppose now that  $s_1 = (i_1, j_1, k) \in H(Q)$  and (4.2) holds for  $s = s_1$ .

**Lemma 4.3.** *One of the following claims holds.*

(i).  $Q_I = \{i_1, i_2, \dots, i_h\}$ , where  $i_1, i_2, \dots, i_h$  are distinct, and  $Q_J \supseteq \{j_1, j_2, \dots, j_\tau\}$ , where  $\tau \leq h$ ,  $j_1, j_2, \dots, j_\tau$  are distinct, such that

$$x(C(s_\nu)) > \frac{1}{2h-1}, \quad (4.3)$$

for  $\nu = 2, \dots, h$ , where  $s_\nu = (i_\nu, j_\nu, k)$ , and  $j_\nu \in \{j_1, \dots, j_\tau\}$  for  $\tau < j \leq h$ ;

(ii).  $Q_I \supseteq \{i_1, i_2, \dots, i_\tau\}$ , where  $\tau \leq h$ ,  $i_1, i_2, \dots, i_\tau$  are distinct, and  $Q_J = \{j_1, j_2, \dots, j_h\}$ , where  $j_1, j_2, \dots, j_h$  are distinct, such that (4.3) holds, for  $\nu = 2, \dots, h$ , where  $s_\nu = (i_\nu, j_\nu, k)$ , and  $i_\nu \in \{i_1, \dots, i_\tau\}$  for  $\tau < i \leq h$ .

*Proof:* Suppose that (4.3) holds for  $s_\nu$ ,  $\nu = 2, \dots, \tau$  such that there are no repetitions in  $\{i_1, \dots, i_\tau\}$  and  $\{j_1, \dots, j_\tau\}$ . Suppose that such a collection is maximal, i.e., for any  $i \in Q_I \setminus \{i_1, \dots, i_\tau\}$  and  $j \in Q_J \setminus \{j_1, \dots, j_\tau\}$ ,

$$x(C((i, j, k))) \leq \frac{1}{2h-1}. \quad (4.4)$$

If the conclusion of this lemma does not hold, then there exists  $i_0 \in Q_I \setminus \{i_1, \dots, i_\tau\}$  and  $j_0 \in Q_J \setminus \{j_1, \dots, j_\tau\}$  such that

$$x(C((i_0, j_\nu, k))) \leq \frac{1}{2h-1} \quad (4.5)$$

$$x(C((i_\nu, j_0, k))) \leq \frac{1}{2h-1}, \quad (4.6)$$

for  $\nu = 1, \dots, \tau$ . However,

$$x(S(Q)) \leq x(S(Q')) + \sum \{x(C(i_0, j, k)) : j \in Q_J, j \neq j_0\} + \sum \{x(C(i, j_0, k)) : i \in Q_I\}, \quad (4.7)$$

where  $Q'$  is defined by  $Q'_I = Q_I \setminus \{i_0\}$ ,  $Q'_J = Q_J \setminus \{j_0\}$ ,  $Q'_K = Q_K = \{k\}$ . By (4.4), (4.5) and (4.6), each term in the right hand side of (4.7), except the first term, is not greater than  $\frac{1}{2h-1}$ . The number of such terms is  $2h-1$ . Clearly,  $Q' \in \bar{S}^{h-1}$ . Thus,  $x(S(Q')) \leq h-1$  by our assumption. Thus, the right hand side of (4.7) is not great than  $h$ . This forms a contradiction with our assumptions on  $x$  and  $Q$ . Therefore, this lemma holds. ■

The following is a brief proof of Theorem 4.1. Cumbersome details are omitted.

*Proof of Theorem 4.1:* By changing 12 to  $4(2h-1)$  in (3.16), and (3.17), we may calculate all  $x(C(s)) > \frac{1}{2h-1}$  in  $O(n^3)$  steps. Check  $s \in S$  with  $x(C(s)) \geq \frac{1}{h}$ . The number of such  $s$  is  $O(n^2)$ . Let  $s_1 = s = (i_1, j_1, k_1)$ . We first check  $Q \in \bar{S}^h$  such that  $s \subseteq Q$  and  $Q$  has a structure as described in Lemma 4.3 (i).

Let  $k = k_1$ . For  $\tau = 2, \dots, h$ , find all possible  $s_\nu = (i_\nu, j_\nu, k)$  for  $\nu = 2, \dots, \tau$  such that (4.3) holds and there are no repetitions in  $\{i_1, \dots, i_\tau\}$  and  $\{j_1, \dots, j_\tau\}$ . (We may find  $s_2, \dots, s_\tau$  sequentially.) Similarly to Lemma 3.4, one may see that the number of  $t = (i', j', k)$  such that

$$x(C(t)) > \frac{1}{2h-1}$$

is  $O(n)$ . Thus, the number of such  $(s_2, \dots, s_\tau)$  is  $O(n^{\tau-1})$ . Then we find all possible  $s_\nu = (i_\nu, j_\nu, k)$  for  $\nu = \tau+1, \dots, h$  such either (i) or (ii) of Lemma 4.3 holds. With an argument similar to Lemma 3.4, one may see that the number of such  $s_\nu$  is  $O(1)$ . In order to expand  $s_1 \cup \dots \cup s_h$  to a set  $Q \in \bar{S}^h$ , we need to add it with

$$\{j_{\tau+1}, \dots, j_h\} \subseteq J \setminus \{j_1, \dots, j_\tau\}.$$



Therefore, the number of expansion ways is  $O(n^{h-\tau})$ . For each  $Q$  produced in this way, we may use formulas similar to (3.6)-(3.8), (3.15) to calculate  $x(S(Q))$  and check whether

$$x(S(Q)) \leq h. \quad (4.8)$$

The complexity of such formulas are  $O(1)$ . Summing up the above discussion, one see that the complexity of check (4.8) for all  $Q \in \bar{S}^h$  such that  $s \subseteq Q$  and  $Q$  has a structure as described in Lemma 4.3 (i), for a fixed  $s$ , is  $O(n^{h-1})$ .

The case of Lemma 4.3 (ii) and the cases, where  $i_1$  or  $j_1$  in  $s$  is fixed, are similar. Since the number of such  $s$  is  $O(n^2)$ , the overall complexity of such a procedure is  $O(n^{h+1})$ . This completes the proof of this theorem. ■

In general, the order of procedures to detect violated facets induced by sets in  $\Phi_h$  for  $h \geq 3$  may be higher than  $O(n^{h+1})$ . For example, let  $h = 4$ . We may find  $O(n^6)$  triples of  $(s, t, p)$  such that  $s, t, p \in S$ ,  $x(C(s)), x(C(t)), x(C(p)) > \frac{1}{4}$ , and

$$a^s \cdot a^t = 0, a^t \cdot a^p = 0, a^s \cdot a^p = 0.$$

Then  $Q := s \cup t \cup p \in S^4$ . We need to check whether

$$x(S(Q)) \leq 4$$

or not. Thus, the complexity of such a procedure is at least  $O(n^6)$ . But  $6 > h+1 \equiv 5$ . Hence, the complexity for detecting violated facets induced by sets in  $\Phi_h$  may be higher than the complexity corresponding to  $\bar{\Phi}_h$  for  $h \geq 3$ .

## REFERENCES

- [1] E. Balas and M. J. Saltzman, "Facets of the three-index assignment polytope", *Discrete Applied Math.* 23 (1989) 201-229.
- [2] E. Balas and M. J. Saltzman, "An algorithm for the three-index assignment problem", to appear in: *Oper. Res.*
- [3] A. M. Frieze, "Complexity of a 3-dimensional assignment problem", *European J. Oper. Res.* 13 (1983) 161-164.
- [4] A. M. Frieze and J. Yadegar, "An algorithm for solving 3-dimensional assignment problem with application to scheduling a teaching practice", *J. Oper. Res. Soc.* 32 (1981) 989-995.

- [5] P. Hansen and L. Kaufman, "A primal-dual algorithm for the three-dimensional assignment problem", *Cahiers du CERO* 15 (1973) 327-336.
- [6] O. Leue, "Methoden zur Lösung dreidimensionaler Zuordnungsprobleme", *Angew. Inform.* (1972) 154-162.
- [7] W. P. Pierskalla, "The tri-substitution method for the three-dimensional assignment problem", *CORS J.* 5 (1967) 71-81.
- [8] W. P. Pierskalla, "The multidimensional assignment problem", *Oper. Res.* 16 (1968) 422-431.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER MSRR-562	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) LINEAR-TIME SEPARATION ALGORITHMS FOR THE THREE- INDEX ASSIGNMENT POLYTOPE		5. TYPE OF REPORT & PERIOD COVERED Technical paper; Sept 1990
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Liqun Qi Egon Balas		8. CONTRACT OR GRANT NUMBER(s) N00014-85-K-0198
9. PERFORMING ORGANIZATION NAME AND ADDRESS Graduate School of Industrial Administration Carnegie Mellon University Pittsburgh, PA 15213		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research (Code 434) Arlington, VA 22217		12. REPORT DATE September 1990
		13. NUMBER OF PAGES 15
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Three-index assignment Facets Cliques Odd holes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Balas and Saltzman (1) identified several classes of facet inducing inequalities for the three-index assignment polytope, and gave $O(n^3)$ separation algorithms for these two classes of facets, and also for a third class. Since the three-index assignment problem has $n^3$ variables, these algorithms are linear-time and their complexity is best possible.		